



**Europäisches
Patentamt**

**European
Patent Office**

**Office européen
des brevets**

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

01123584.3

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk



Anmeldung Nr.:
Application no.: 01123584.3
Demande no:

Anmeldetag:
Date of filing: 01.10.01
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

SOCIETE DES PRODUITS NESTLE S.A.
Case postale 353
1800 Vevey
SUISSE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se referer à la description.)

Cocoa flavour precursor peptides

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

C07K/

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE TR

01. Okt. 2001

Cocoa flavour precursor peptides

5

The present invention pertains to peptides derived from fermented cocoa beans and representing cocoa and/or chocolate flavour precursors. In particular, the present invention relates to a process for preparing cocoa and/or chocolate flavour precursor peptides from fermented cocoa beans and to the preparation of cocoa/chocolate flavour on a synthetic basis, and to the use thereof in the preparation of chocolate.

10

The processing of cocoa beans comprises the basic steps of fermentation, drying and roasting.

15 During fermentation the cocoa seed proteins are degraded by microbiological and enzymatic processes to mainly hydrophilic peptides and hydrophobic amino acids, which serve as flavour precursors for the unique chocolate flavour. The cocoa proteolytic system involved is highly efficient. The endogenous proteolytic activity in cocoa is represented by endo-
proteases, aminopeptidases and carboxypeptidases, which in concert have the capability to
20 hydrolyze intact protein to single amino acids and oligopeptides of different length.

20

During roasting various chemical reactions occur, in particular Maillard-type reactions and thermal degradation reactions (decarboxylations, deaminations, oxidations etc.). The Maillard reaction involving peptides, amino acids and reducing sugars generates compounds
25 that are deemed to eventually contribute to the final cocoa/chocolate flavour.

25

In the recent past more and more evidence has been brought up that the size of the peptides and their amino acid contents/sequence play an important role in flavour formation. The amino acid pool in un-roasted and roasted fermented cocoa has been investigated and certain
30 hydrophobic amino acids have been found to be implicated in the generation of cocoa flavour (Seiki, Rev. Int. Choc. 28 (1973) 38-42). In addition several attempts to produce artificially

30

cocoa flavour are known in the art, such as e.g. subjecting acetone dried powder prepared from unfermented ripe cocoa beans to an autolysis at a pH of 5.2 followed by roasting in the presence of reducing sugars (Voigt et al., Food Chem. 49 (1994), 173-180).

5 However, little is known in the art about short-chain peptides present in the cocoa peptide pool. So far, merely four major proteins with an apparent molecular weight of 14.5, 21, 31 and 47 kDa, respectively, could be identified in cocoa beans before fermentation, which proteins are deemed to be the main source for the peptide/amino acid pool, generated during fermentation.

10

One of the major limitations in studying the peptide pool in fermented cocoa beans is the presence of UV absorbing compounds, e.g. theobromine, caffeine, epicatechin, or polyphenols in general. Extraction of the proteinaceous content in the presence of such compounds generally leads to extensive complexation. As a consequence, trying to remove
15 said polyphenols with current solvent systems, such as aqueous acetone or methanol, was not successful, as these solvent systems also remove low-molecular-peptides.

Since so far no substantial information about short-chain peptides present in fermented cocoa beans is presently available an object of the present invention resides in providing more
20 detailed data about said peptides and eventually provide means for improving the preparation of cocoa flavour.

The above object has been solved by providing peptides obtainable by a process comprising the steps of

- 25 (i) preparing a cocoa nib powder starting from fermented cocoa beans;
(ii) extracting said cocoa nib powder with aqueous acetic acid (50 %);
(iii) separating non-proteinaceous compounds with a solid phase adsorption and collecting the elute containing peptides;
(iv) diluting the elute with 5 volumes of 0.1 % trifluoroacetic acid;
30 (v) loading the diluted elute on a RP-HPLC column equilibrated with 0.14 % sodium acetate/0.05 % TEA and eluting with an increasing concentration of 80

% acetonitrile/0.1 % trifluoroacetic acid;

(vi) collecting the elutes containing the peptides

5 The peptides according to the present invention have been shown to exhibit a size of approximately 2 to 25 amino acids and may serve as precursors for flavour active substances.

In the figures,

10 Fig. 1 shows an evaluation of different solvents to extract proteinaceous material from fermented cocoa beans;

Fig. 2 shows the results of RP-HPLC purification as determined by UV absorbance (A) and a fluorimetric assay (B,C).

15 During the extensive studies leading to the invention, the present inventors have designed novel methods for the isolation of short-chain cocoa peptides derived from fermented cocoa beans.

20 The peptides or peptide pool derived from fermented cocoa beans of the present invention are obtainable by a process comprising the above steps (i) to (iv).

During step (i) fermented cocoa beans are crushed by conventional means, such as a universal mill with the grinded material being preferably sieved to remove remaining large particles.

25 In order to be capable to primarily extract the peptides of interest extensive studies have been carried out to find an appropriate solvent system. To this end, several solvent systems have been investigated involving citrate phosphate buffer (50mM; pH 3), citrate phosphate buffer (50mM; pH 8), 50 mM sodium phosphate buffer, pH 8 plus 0.5 M NaCl, methanol, 10 mM
30 potassium phosphate buffer, pH 7.0, isopropanol, formic acid and acetic acid (Fig. 1).

Acetic acid (50%) provided an overall recovery of total amino groups of about 70 % with the additional advantage of its easy removal under low pressure and its ability to inhibit oxidation reactions. Moreover, it has been found that by using this particular solvent there is no need to carry out laborious and time consuming processes of fat removal with lipophilic solvents prior to extraction, such that both intact or defatted cocoa nib powder may be used.

In order to provide best results the cocoa nib powder is suspended in the solvent, sonicated and optionally further homogenized by appropriate means. The suspension may be cooled on ice and centrifuged at 15.000 rpm at 4 °C for 15 min to obtain a clear supernatant between the pellet and the fat layer at the surface. For further purification this supernatant may be separated, passed through a 0.22 µm filter disc and optionally kept at -20 °C until further processing.

During step (iii) the acetic acid extract obtained is subjected to a treatment to remove non-proteinaceous, organic compounds, in particular UV-absorbing compounds and polyphenols, respectively. This may be achieved by means of a solid phase extraction with subsequent centrifugation of the elute and isolation of the supernatant solution. To this end a Chromabond® cartridge (obtainable from Macherey-Nagel) equilibrated with 0.1 % TFA (trifluoroacetic acid) may be used on which the extract is loaded. For differentially eluting the adsorbed material from the column a solvent system consisting of 70 % methanol in 0.1 % trifluoroacetic acid may be used. However, also other solvent systems may be put to use with the results of the elution being monitored via conventional means comprising mass spectrometry, UV/VIS or fluorometric assays in combination with UV/VIS.

The respective elute(s) comprising the peptides of interest is(are) then dried under reduced pressure to remove the solvent and is subsequently dissolved in an appropriate solvent for further processing, such as 0.1 % TFA or 25 % acetonitrile/0.1% TFA (v/v).

Following, the solution obtained as above is diluted with about 5 volumes of 0.1 % trifluoroacetic acid and is loaded on a RP-HPLC column equilibrated with 0.14 % sodium acetate/0.05 % TEA. The adsorbed material is eluted with increasing concentrations of a

solvent comprising 80 % acetonitrile/0.1 % trifluoroacetic acid.

Finally, the elutes containing the objective peptides are collected and optionally dried and dissolved in an appropriate solvent.

5

The peptides obtainable according to the above described methods have a size of between about 2 to 25 amino acids. In order to obtain compounds exhibiting a cocoa and/or chocolate flavour, these peptides obtained as above may be subjected to a Maillard reaction involving reducing sugars or may be subjected to a further enzymatic degradation using e.g. aspartic endoproteinase, cysteine endoproteinase, carboxypeptidase. Consequently, the peptides of the present invention or the products or fragments thereof may serve as precursors of flavour-active compounds. The flavour-active substances obtainable according to the present invention may be added to food products, cosmetic products and/or pharmaceutical products.

10

15 The present invention also pertains to a process for obtaining cocoa/chocolate flavour precursors, which process comprises the steps (i) to (vi) as detailed above.

The following examples illustrate the invention in a more detailed manner. It is, however, understood that the present invention is not limited to the examples but is rather embraced by the scope of the appended claims.

20

Example 1

Preparation of fermented cocoa nib powder

25

West African Amelonado cocoa beans were fermented in 80 kg heaps covered with banana leaves in Ivory Coast. Samples were removed at 1 day intervals during 7 day fermentation period, sun dried, hand peeled and shipped. The cocoa beans were kept at -20 °C.

30 Dried cocoa beans were passed through a bean crusher followed by a winnover to remove shells. Cocoa nibs were milled for a few seconds in an universal mill. The nib powder was passed through 0.8-mm sieve and kept at 4 °C. Unless stated otherwise, the fermented beans

used throughout the present study were 5 day Amelonado beans.

Preparation of unfermented cocoa nib powder

- 5 Cocoa nib powder was prepared from unfermented cocoa beans in an analogous manner.

Extraction of Cocoa nib powder

Cocoa nib powder (2 g) was suspended in 20 ml of a solvent system [citrate phosphate buffer (50mM; pH 3), citrate phosphate buffer (50mM; pH 8), 50 mM sodium phosphate buffer, pH 8 plus 0.5 M NaCl, methanol, 10 mM potassium phosphate buffer, pH 7.0, isopropanol, formic acid (70 %) and acetic acid (50 %)]. The resulting suspension was sonicated for 1 min at full speed (10 mm probe at medium setting and full power, Labsonic U, B. Braun). The suspension was then further subjected to homogenization treatment for 1 min at a maximum speed (Polytron, Kinematica AG). The suspension was cooled on ice for 15-30 min and centrifuged at 15.000 rpm at 4 °C for 15 min. The clear supernatant located between the pellet and the fat layer at the surface was carefully removed, passed through 0.22 µm filter disc and kept at -20 °C until further processing.

- 20 In order to determine the effect of the extraction procedure the amount of amino groups present in the sample of the fermented CNP as well as in the extracts were determined by acid hydrolysis (6N HCl at reduced pressure at 110 °C for 24 hrs).

To this end, aliquots of the respective samples (10 µl) were dried under reduced pressure in Pyrex glass tubes and subsequently acid hydrolyzed under vapor phase of HCl using Waters Pico Tag Workstation equipped with Savant SpeedVac concentrator (refrigerated trap RVT 400; SVC 100 H centrifuge and Alc-Tel vacuum pump). Samples were flushed with nitrogen followed by applying reduced pressure until few air bubbles were visually detected. The process was repeated 5-times and the tubes were sealed under reduced pressure and placed at 110 °C for 24 hrs. After hydrolysis, samples were brought to room temperature and the vacuum applied was carefully bled, to avoid splashing of HCl. The excess HCl was removed

and the hydrolyzed samples were dried under reduced pressure after the addition of 20 μ l of re-dry buffer (water/methanol/TEA (triethylamine); 2:2:1) and suspended in 200 μ l of water.

The hydrolyzed samples (10 μ l) were dried in a SpeedVac and dissolved in 10 μ l re-dry
5 buffer and again dried. The dried samples were dissolved in 20 μ l of derivatization solution (ethanol/water/TEA/PITC (phenylisothiocyanate); 7:1:1:1), vortexed and incubated at room temperature for 10 minutes. The samples were dried in a rotary SpeedVac for 60 minutes (reduced pressure 50-60 m Torr) and dissolved in 100 μ l solvent A (0.14 M Na-acetate/0.05% TEA) and subjected to RP-HPLC for PITC derivatized amino acid analysis
10 (Heirikson and Meredith Anal. Biochem. 136 (1984), 65-74).

The RP-HPLC was carried out on Hewlett Packard 1050 HPLC system, with the detection being carried out with an UV/VIS diode array detector. The solvents were degassed on-line using Gastorr G-104 automatic degassers. The derivatized amino acids (10 μ l) were injected
15 on to reverse-phase C18 column (AccQ.Tag column (3.9x150mm) from Waters) and the amino acid derivatives were eluted with an increasing linear gradient of solvent B (60 % v/v acetonitrile) in solvent A (supra) as follows: 0-48 % B in 10 min; 48 - 100 % in 0.2 min followed by isocratic elution at 100 % B for 1.5 min. Peaks were recorded and integrated by HP chemstation software.

20 It could be shown that intact proteins and peptides were best dissolved in acetic acid and formic acid, which gave a 2-fold better recovery as compared to the other solvent systems. Due to its ease of handling acetic acid has been chosen.

25 Example 2

Separation of the amino acid/peptide pool from polyphenols

In order to separate the short-chain peptides from other organic compounds solid phase extraction employing Chromabond® C8 cartridge (500 mg) has been utilized. Acetic acid
30 extracts as obtained in example 1 were diluted 5-times with water, chilled on ice for 15 min and centrifuged (5.000 rpm) for 15 min at 4 °C yielding a clear supernatant. Chromabond®

cartridge was pre-conditioned by passing 1 ml of methanol followed by 2 ml 50 % methanol in 0.1 % TFA. The cartridge was equilibrated with 3 ml 0.1 % TFA. The sample (1 ml) was passed through the column slowly followed by 2-times 1 ml wash with 10 % methanol in 0.1 % TFA. The adsorbed material from the column was eluted with 2-times 1 ml of 70 % methanol on 0.1 % TFA. The column wash and 10 % methanol elute fractions were pooled (SPE elute A), dried under low pressure and redissolved in 0.1 % TFA. The 70 % elute fraction (SPE elute B) was also dried under low pressure and redissolved in 25 % ACN (acetonitrile)/0.1 % TFA.

- 10 The determination of the total primary amino groups and amino acid analysis before and after acid hydrolysis (supra, example 1) showed that SPE elute A represents over 90 % of the total soluble proteinaceous material (Table 1). SPE elute A from acetic acid extract of fermented beans showed a 3-fold increase following acid hydrolysis indicating presence of peptides (Table 1). The average chain length of peptides was 3 amino acid residues.

Table 1

	Amino groups		Recovery, %	
	N_{α}^a	N_t^b	N_{α}^a	N_t^b
	<i>μmol</i>			
Unfermented				
Supernatant	35	38	100	100
SPE elute A	29	26	83	68
SPE elute B	1	6	3	16
Fermented				
Supernatant	90	218	100	100
SPE elute A	68	127	76	58
SPE elute B	11	81	12	37

^a Before and ^b after acid hydrolysis (6N HCl at 110 °C for 24 h). N_{α} and N_t , Free and total amino groups.

Example 3

Separation of peptides by HPLC

Short-chain hydrophilic peptides and the majority of amino acids are in general not retained
5 on a reverse-phase HPLC column due to their poor interactions with the hydrophobic chains
(C8 or C18) of the stationary phase. The interactions of hydrophilic peptides and amino acids
can even be improved by the inclusion of aliphatic sulfonic acids, e.g. hexane- or heptane-
sulfonic acid. In the case of ion-pair RP-HPLC, 5 mM sodium hexane sulfonate was included
in solvent A and B.

10 Samples (20 μ l) were injected onto the RP-column (Spherisorb 80-5C8 (250 x 4.6 mm)) and
eluted with a solvent system comprising solvent A (0.1 % v/v TFA in water), to which
hexanesulfonic acid has been added (5mM) and solvent B (80 % v/v CAN and 0.1 % v/v
TFA), also including hexanesulfonic acid in the same amount, with isocratic at 0 % solvent B
15 for 10 minutes, 0-50 % B in 60 minutes, isocratic at 50 % B for 5 min, 50-100 % B in 25
minutes and isocratic at 100 % B for 5 minutes.

The peaks were detected by UV absorbance at 205 nm (fig. 2A) and by fluorometric
detection following post column reaction with a o-phthaldialdehyde (OPA) reagent (Fig. 2B)
20 (Jarrett et al. Anal. Biochem. 153 (1986), 189-198).

An optimized gradient (0 % B for 20 min, 0-25 % B in 70 min; 25 - 100 % B in 10 min; and
isocratic elution at 100 % B for 10 min) gave best results as may be seen in Fig. 2C.

25 The same assay as above was repeated without the addition of hexanesulfonic acid and the
elutes during the first 10 minutes were analyzed by electrospray ionization (ESI) mass
spectrometry using a FinniganMat LCQ mass spectrometer interfaced with an HPLC (Spectra
system, FinniganMat). The HPLC system (Spectra system, FinniganMat) interfaced with
LCQ mass spectrometer (FinniganMat) consisted of a quaternary pump (TSP P4000), an
30 autoinjector (TSP AS3000) and a UV/VIS detector (UVIS 205 from Linear Instruments)
equipped with high pressure stainless steel flow cell (1.6 μ l volume, 2 mm path-length). LC

flow was directed to LCQ mass spectrometer without using a flow-splitter. Typically the following conditions were used: capillary temperature, 200°C, sheath gas flow, 70; auxiliary gas flow, 10; source voltage, 5kV. Other parameters were adjusted automatically during calibration /tuning procedure as recommended by the manufacture.

5

A comparison of molecular ions data $[(M+H)]^+$ from fermented and non fermented cocoa beans showed the presence of 43 additional molecular ions in the fermented beans elute.

10

Each identified molecular mass ion was upon MS/MS analysis, and in conjunction with on-line data analysis using Sequest™ database, identified various peptides- both of cocoa vicillin and albumin origin. 18 peptide sequences from vicillin and 25 peptide sequences from albumin, a total of 43 sequences, were observed.

Table 3
Masses of peptides deduced using SequestTM

Peptide, <i>m/z</i>	Molecular ion	Sequence	Source protein
698.4		PVNSPGK	Vicilin
861.4		PVNSPGKY	Vicilin
1118.5		NADSKDDVVR	Albumin
1205.6		SNADSKDDVVR	Albumin
1391.6		SNADSKDDVVRVS	Albumin
629.3		VPHYN	Vicilin
515.3		VPHY	Vicilin
784.4		ENSPPLK	Vicilin
855.4		AENSPPLK	Vicilin
727.4		AENSPPL	Vicilin
600.3		KDQPL	Vicilin
872.4		ASKDQPLN	Vicilin
758.4		ASKDQPL	Vicilin
487.2		KAGVL	Albumin
544.3		KAGVLG	Albumin
857.5		KIEKAGVL	Albumin
431.2		AGGGGL	Albumin
688.3		ISGAGGGGL	Albumin
933.4		SSISGAGGGGLA	Albumin
862.5		SSISGAGGGGL	Albumin
838.4		DEEGNFK	Vicilin
373.3		KIL	Vicilin
710.3		DEEGNF	Vicilin
300.2		APL	Vicilin
621.3		SPGDVF	Vicilin
902.5		APLSPGDVF	Vicilin
826.4		RNNPYY	Vicilin
1042.5		NNPYYPFK	Vicilin
940.4		YDNSAGKW	Albumin
1054.5		NYDNSAGKW	Albumin
892.4		DTDGDELQ	Albumin
1050.4		DTDGDELQTG	Albumin
715.4		ANSPVLD	Albumin
600.3		ANSPVL	Albumin
563.3		DNEW	Albumin
820.3		DNEWAW	Albumin
475.3		PVIF	Albumin
576.3		TPVIF	Albumin
747.4		NGTPVIF	Albumin
555.3		AGHAVT	Vicilin
453.2		DYR	Vicilin
690.4		NGKGTIT	Vicilin
403.2		DRL	Albumin
484.3		VPIR	Albumin
822.4		SNQNGRF	Vicilin
460.3		DIGR	Albumin
634.3		VSTDVN	Albumin
453.2		DYR	Vicilin
1367.9		PAGNNKPESYYGA	Vicilin
328.2		VPI	Albumin
435.3		RIF	Vicilin
784.4		GEPGPNTL	Albumin

Example 4

Roasting experiments

- 5 In a subsequent step, it was determined which peptides and amino acids are potentially yielding flavour active substances, when subjected to a reaction during the roasting step.

To this end, fermented and dried cocoa beans were roasted at 145 °C for 30 min and in addition one sample for each day of fermentation was prepared. The roasted beans were
10 ground and extraction was carried out. For comparison, non-roasted beans were processed at the same time as roasted beans. LC-MS/MS carried out on both roasted and non-roasted beans.

As a result, it was observed that nearly all peptides from vicilin and albumin fragmentation
15 appear after roasting, but a substantial reduction in their quantity could be observed.

Claims

01. Okt. 2001

1. Cocoa and/or chocolate flavour precursor peptides obtainable by a process comprising the steps of:

- 5 (i) preparing a cocoa nib powder starting from fermented cocoa beans;
(ii) extracting said cocoa nib powder with aqueous acetic acid (50 %);
(iii) separating non-proteinaceous compounds with a solid phase adsorption and collecting the elute containing peptides;
(iv) diluting the elute with 5 volumes of 0.1 % trifluoroacetic acid;
10 (v) loading the elute fraction on a RP-HPLC column equilibrated with 0.14 % sodium acetate/0.05 % TEA and eluting with an increasing concentration of 80 % acetonitrile/0.1 % trifluoroacetic acid;
(vi) collecting the elutes containing the peptides

- 15 2. The peptides according to claim 1, wherein the peptides have a size of about 2 to 30 amino acids, preferably 2 to 5 amino acids.

3. A process for preparing cocoa and/or chocolate flavour, which comprises, subjecting the peptides obtained according to claim 1 or 2 to a Maillard reaction with reducing sugars.

- 20 4. The process according to claim 3, which comprises the step of further digesting the peptides obtained according to a process according to claim 1 with proteases.

5. Use of a peptide according to any of the claims 1 or 2 for the preparation of cocoa and/or chocolate flavour.

- 25 7. Use of a peptide according to any of the claims 1 or 2 for the preparation of confectionery, ice cream, beverages, dairy products, cosmetic products, pet food or pharmaceutical products.

- 30 8. The use according to claim 7, wherein the confectionery product is chocolate.

Summary

5 The present invention pertains to peptides derived from fermented cocoa beans and representing cocoa and/or chocolate flavour precursors. In particular, the present invention relates to a process for preparing cocoa and/or chocolate flavour precursor peptides from fermented cocoa beans and to the preparation of cocoa/chocolate flavour on a synthetic basis, and to the use thereof in the preparation of chocolate.

01. Okt. 2001




Fig.1 Evaluation of different solvents to extract cocoa proteinaceous material. The per cent of solvent indicated is by vol. Unless indicated otherwise each extraction was carried out at concentrations of 10 % (w/v). Total amino groups and amino acids in the acid hydrolyzed cocoa nib powder were assumed to be maximum extractable amounts. (■) N_t, total amino groups and (▨) A_t, total amino acids. For details see "Experimental Procedures". CPB3, citrate phosphate buffer (50 mM; pH 3), CPB8, citrate phosphate buffer (50 mM; pH 8), PB8/NaCl, 50 mM sodium phosphate buffer, pH 8 plus 0.5 M NaCl, MeOH, methanol, PB/7, 10 mM potassium phosphate buffer, pH 7.0, ISP, isopropanol, FA, formic acid, AcA, acetic acid.

BEST AVAILABLE COPY




Fig. 2 RP-HPLC of SPE fractions. Appropriately diluted samples were injected onto RP column (Spherisorb 80-5C8 (250 x 4.6 mm)) and eluted with TFA/HSA/ACN solvent system (see Experimental Procedure for) with gradient isocratic at 0 % B for 10 min, 0-50 % B in 60 min, isocratic at 40 % B for 5 min, 50-100 % B in 25 min and isocratic at 100 % B for 5 min. The peaks were detected by UV absorbance at 205 nm (A) and by fluorometric detection (B) following post-column reaction with OPA reagent. C, Optimized gradient (0 % B for 20 min, 0-25 % B in 70 min, 25 to 100 % B in 10 min and isocratic elution at 100 % B for 10 min. Red trace, fermented AcA extract and blue trace, unfermented AcA extract